



^{*}Where possible information should be provided as a leaflet (leaflets can be ordered here).

^{*}RAVS entry is a live-system that auto populates the GP record.

London opportunistic maternity immunisation offer pathway (General Practice)





Box 2: Immunisations delivery timelines

Pertussis can be administered from 16 weeks up to 8 weeks after birth

RSV can be administered from 28 weeks up to birth


Influenza can be administered at any point in the pregnancy during flu season (1st September to 31st March) and outlined in the annual flu letter)

Key information for primary care

- Ensure you have sufficient vaccine stock to enable opportunistic offers
- Ensure appropriate immunisation information is provided around all vaccinations at all visits
 - Where possible immunisation information should be provided as a leaflet (leaflets can be ordered <u>here</u>).

Terminology

The term woman is used in this document to encompass all pregnant people including pregnant women, trans men, and those who identify as non-binary.



^{*}Where possible immunisation information should be provided as a leaflet (leaflets can be ordered here).

^{**}Separate immunisation appointment/(s) with practice nurse may need to be booked. See co-administration guidance in Appendix.

Appendix

- Co administration of RSV with Pertussis: NHS England » General practice respiratory syncytial virus (RSV) vaccination programme: contractual guidance
- Green Book chapters:
 - Respiratory syncytial virus: the green book, chapter 27a GOV.UK
 - Pertussis: the green book, chapter 24 GOV.UK
 - Influenza: the green book, chapter 19 GOV.UK
 - COVID-19: the green book, chapter 14a GOV.UK