

Remember...

Some inhalers must be used with a spacer.
Check with your GP, asthma nurse or pharmacist

Always keep your rescue inhaler and your spacer with you. You might need them if your asthma gets worse

Make sure you have an asthma review within 48 hours after an attack

Vaping in children and young people is dangerous and can cause permanent lung damage

My Asthma Triggers

List the things that make your asthma worse:

Pollen	Vaping
Dust	Environmental pollution
Animal fur	Other fumes/ sprays
Weather	Respiratory infections (cold/flu)
Exercise	Medicines
Mould/damp	Stress/emotions
Fumes	Food *
Tobacco smoke	
House dust mite	

* Always refer to your Allergy Plan as well

Any Other Triggers:

REMEMBER

Good asthma control means having **NO** symptoms

If you have any symptoms you should speak to your doctor or asthma specialist as soon as possible



MART Asthma Plan

Ages 12 - 18

Name:
Date:

Produced by
London Babies, Children and Young People's Team

Extra Advice from my Asthma Professional:

Contact Details

GP:

Asthma Specialist/Team:

Additional Resources:

[Asthma and Lung UK
Asthma Toolkit](#)



[Check you're using
your inhaler properly:](#)



This plan was approved by
London Asthma Leadership and Implementation Group (LALIG)
Approval Date: Jan 2025
To Be Revised: Jan 2027

Every day:
I am symptom free



I only use one inhaler for my asthma

For maintenance

I need to take my inhaler every day for maintenance

It is called:

and its colour is:

My best peakflow measure is: l/min

I take puff/s of my maintenance inhaler in the morning and puff/s at night. I do this every day even if my asthma's OK.

Other asthma medicines I take every day:

For rescue

When I am having symptoms, I will use the same inhaler. I should not need rescue puffs regularly.

I take puff/s when needed.

My asthma is not
controlled if...



I wheeze, cough, my chest hurts, or it's hard to breathe **or**

If my asthma is stopping me doing sport or other activity **or**

I'm waking up at night because of my asthma **or**

My peakflow measure falls below 80%:

l/min

So I need to...

Take puff/s of my inhaler.

After 2 minutes if I still have symptoms, repeat this, one puff at a time, until my symptoms improve. I can take up to puffs.

My symptoms should be gone. If they come back I can repeat this process.

I will arrange
to see my GP
today or
tomorrow if...

I need to take rescue puffs more than two days a week **or** more than once in 24 hours.

I will go to the red
section if...

My asthma symptoms are not improving even with the extra rescue doses **or**

My symptoms come back quickly **or**

My symptoms have come back but I have taken my maximum daily puffs (puffs).

I'm having an asthma
attack and need urgent
attention and...



I will see a doctor
urgently
(emergency
department or GP
if available) if...

My symptoms have come back but I have taken my maximum daily puffs **or**

My symptoms come back quickly **or**

My peakflow measure falls below 60%:
 l/min

I will call 999
and tell them
I'm having an
asthma attack
if...

If my symptoms aren't completely better after puffs in a row **or**

My symptoms are getting worse.

If I still have symptoms, I can continue to take additional puffs as required.

I also need to...

Sit up - don't lie down. Try to keep calm.

If the ambulance has not arrived after 10 minutes, **contact 999 again immediately**, and take additional puffs every 2 minutes if I still have symptoms.

If I don't have my MART inhaler with me and need to use the emergency blue reliever inhaler, take one dose every 30-60 seconds up to a maximum of 10 puffs.