

Pharmacy First – examples of conditions general practice can refer to community pharmacy for minor illness

North East London

Pharmacy First: minor illness consultations

This is not an exhaustive list, Community pharmacies can receive referrals for anything considered a minor illness.

Acne, spots, and pimples	Allergic reaction	Ankle or foot pain or swelling	Athlete's foot	Bites or stings, insect, or spider	Blisters
Constipation	Cough	Cold and flu	Diarrhoea	Ear discharge or ear wax	Earache
Eye, red or irritable	Eye, sticky or watery	Eyelid problems	Hair loss	Headache	Hearing problems or blocked ear
Hip, thigh, or buttock pain or swelling itch	Knee or lower leg pain	Lower back pain	Lower limb pain or swelling	Mouth ulcers	Nasal congestion
Pain and/or Frequency passing urine	Rectal pain	Scabies	Scratches and grazes	Shoulder pain	Skin, blisters or rash
Sleep difficulties	Sore throat	Teething	Tiredness	Toe pain or swelling	Vaginal discharge
Vaginal itch or soreness	Vomiting	Wound problems – management of dressings	Wrist, hand, or finger pain or swelling		

Pharmacy First - Clinical Pathway (7 clinical conditions)

Please note these are the main exclusions to support referrals. The community pharmacist will review specific symptoms during the consultation

Urinary tract infection	Shingles*	Impetigo	Infected insect bites	Acute sore throat	Acute sinusitis	Acute otitis media
A UTI is an infection in any part of the urinary system.	Shingles is an infection that causes a painful rash	Impetigo is a common infection of the skin. It is contagious, which means it can be passed on by touching.	Insect bites and stings can become infected or cause a reaction.	Sore throat is a symptom resulting from inflammation of the upper respiratory tract	Sinusitis is swelling of the sinuses, usually caused by an infection. The sinuses are small, empty spaces behind your cheekbones and forehead that connect to the inside of the nose.	An infection of the middle ear.
Inclusion: <ul style="list-style-type: none"> Female Aged between 16 - 64 Suspected lower UTI 	Inclusion: <ul style="list-style-type: none"> 18 years and over Suspected case of shingles. Rash appeared within the last 72 hours - 7 days 	Inclusion: <ul style="list-style-type: none"> 1 year and over Signs and symptoms of impetigo Localised (4 or fewer lesions/clusters present) 	Inclusion: <ul style="list-style-type: none"> 1 year and over Infection that is present or worsening at least 48 hours after the initial bite(s) or sting(s) 	Inclusion: <ul style="list-style-type: none"> 5 years and over Suspected sore throat 	Inclusion: <ul style="list-style-type: none"> 12 years and over Suspected signs and symptoms of sinusitis Symptom duration of 10 days or more 	Inclusion: <ul style="list-style-type: none"> Aged between 1 – 17 Suspected signs and symptoms of acute otitis media
Exclusion: <ul style="list-style-type: none"> Male <16 or >64 Pregnant Breastfeeding Recurrent UTI (2 in last 6 months or 3 in last 12 months) Catheter Type 1 or 2 Diabetic 	Exclusion: <ul style="list-style-type: none"> < under the age of 18 Pregnant or suspected pregnancy Breastfeeding with shingle sores on the breasts Shingles rash onset over 7 days ago 	Exclusion: <ul style="list-style-type: none"> < under 1 year of age Pregnancy or suspected pregnancy in individuals under 16 years of age Breastfeeding with impetigo lesion(s) present on the breast Recurrent impetigo (2 or more episodes in the same year) Widespread lesions/ clusters present Systemically unwell 	Exclusion: <ul style="list-style-type: none"> < under 1 year of age Pregnancy or suspected pregnancy in individuals under 16 years of age Systemically unwell Bite or sting occurred while travelling outside the UK 	Exclusion: <ul style="list-style-type: none"> Individuals under 5 years of age Pregnancy or suspected pregnancy in individuals under 16 years of age age Recurrent sore throat/tonsillitis (7 or more significant episodes in the preceding 12 months or 5+ in each of the preceding 2 years, or 3+ in the preceding three years) Previous tonsillectomy 	Exclusion: <ul style="list-style-type: none"> Individuals under 12 years of age Pregnancy or suspected pregnancy in individuals under 16 years of age Symptom duration of less than 10 days Recurrent sinusitis ((4 or more annual episodes of sinusitis) 	Exclusion: <ul style="list-style-type: none"> Individuals under 1 year of age or over 18 years of age Pregnancy or suspected pregnancy in individuals under 16 Recurrent infection (3+ episodes in preceding 6 months, or 4+ episodes in the preceding 12 months with at least one episode in the past 6 months.)